

example of the second-generation neonicotinoids. CGA 293'343 provides excellent control of sucking and chewing insects. Control of most insect pests with CGA 293'343 is superior or equivalent to that of currently registered neonicotinoid insecticides. Structure–activity relationships revealed that variation of the pharmacophore, the oxadiazinane ring, the heterocyclic group and the substituent R in CGA 293'343 diminish biological activity against *Aphis craccivora*.

## REFERENCES

- 1 Yamamoto I, Neonicotinoids – mode of action and selectivity. *Agrochem. Jpn* **68**:14–15 (1996).
- 2 Tomizawa M and Yamamoto I, Binding of nicotinoids and related compounds to the insect nicotinic acetylcholine receptor. *Nihon Noyaku Gakkaishi (J Pestic Sci)* **17**:231–236 (1992).
- 3 Elbert A, Overbeck H, Iwaya K and Tsuboi S, Imidacloprid, a novel systemic nitromethylene analogue insecticide for crop protection. *Proc Brighton Crop Prot Conf – Pests and Diseases*. Vol 1, pp 21–28 (1990).
- 4 Minamida I, Iwanaga K, Tabuchi T *et al*, Synthesis and insecticidal activity of acyclic nitroethene compounds containing a heteroarylethylamino group. *Nihon Noyaku Gakkaishi (J Pestic Sci)* **18**:41–48 (1993).
- 5 Takahashi H, Mitsui J, Takakusa N *et al*, NI-25, a new type of systemic and broad-spectrum insecticide. *Proc Brighton Crop Prot Conf – Pests and Diseases*. vol 1, pp 89–96 (1992).
- 6 Gsell L, Preparation and testing of [(pyridylmethyl)-amino]nitroethylenes as insecticides, acaricides, and ectoparasiticides. *Eur Pat Appl EP 302833 A2 890208* (1987)
- 7 Gsell L, Preparation of pyridylmethylcyanoguanidines as insecticides and acaricides. *Eur Pat Appl EP 306696 A1 890315* (1987).
- 8 Kristiansen O, Maienfisch P and Gsell L, Guanidine derivatives as insecticides. *Eur Pat Appl EP 418199 A2 910320* (1989)
- 9 Maienfisch P, Gonda J, Jacob O *et al*, Synthesis and insecticidal activity of novel nicotinoids. *Book of Abstracts, 214th ACS National Meeting, Las Vegas, NV, 7–11 September AGRO-018*. American Chemical Society, Washington, DC (1997).
- 10 Maienfisch P and Gsell L, Preparation of 3-(heterocyclylmethyl)-4-iminoperhydro-1,3,5-oxadiazine derivatives as pesticides. *Eur Pat Appl EP 580553 A2 940126* (1992).
- 11 McKay AF and Wright GF, Preparation and properties of N-methyl-N'-nitroso-N'-nitroguanidine. *J Am Chem Soc* **69**:3028–3030 (1947).
- 12 Uneme H, Higuchi N and Minamida I, Preparation of chlorothiazole derivatives. *Eur. Pat. Appl., EP 446913 A1 910918* (1990).
- 13 Maienfisch P and Gsell L, Preparation of substituted 4-nitroiminoperhydro-1,3,5-oxadiazine derivatives and their use as pesticides. *PCT Int Appl, WO 9806710 A1 980219* (1996).

## Synthetic approaches towards CGA 293'343: A novel broad-spectrum insecticide

Thomas Göbel,<sup>1</sup> Laurenz Gsell,<sup>2</sup> Ottmar F Hüter,<sup>2</sup> Peter Maienfisch,<sup>2</sup> Rudolf Naef,<sup>3</sup> Anthony C O'Sullivan,<sup>2</sup> Thomas Pittner,<sup>2\*</sup> Thomas Rapold,<sup>4</sup> Gottfried Seifert,<sup>4</sup> Marcel Senn,<sup>5</sup> Henry Szczepanski<sup>2</sup> and David J Wadsworth<sup>1</sup>

<sup>1</sup> Novartis Animal Health Inc, PO Box, CH-4002 Basel, Switzerland

<sup>2</sup> Novartis Crop Protection AG, PO Box, CH-4002 Basel, Switzerland

<sup>3</sup> Novartis Services AG, PO Box, CH-4002 Basel, Switzerland

<sup>4</sup> Novartis Crop Protection Münchwilen AG PO Box, CH-4333 Münchwilen, Switzerland

<sup>5</sup> Novartis Crop Protection Monthey SA, Route de l'Île au Bois CH-1870 Monthey, Switzerland

**Abstract:** Synthetic approaches towards CGA 293'343 (ISO draft common name: thiamethoxam), a novel broad-spectrum insecticide from the class neonicotinoids, are described. 2-Chloro-5-chloromethylthiazole, an important synthetic intermediate, was prepared from five different precursors. Alternatively, CGA 293'343 was prepared via the intermediate 2-benzylmercapto-5-chloromethylthiazole, the synthesis of which is also described.

© 1999 Society of Chemical Industry


**Keywords:** CGA 293'343; thiamethoxam; neonicotinoid; insecticide; thiazole; oxadiazinane; synthesis

## 1 INTRODUCTION

CGA 293'343<sup>1</sup> (Fig 1; ISO draft common name: thiamethoxam), which is a novel broad-spectrum insecticide, shows outstanding activity against a wide spectrum of important pests; it belongs to the class of neonicotinoids<sup>2</sup> and is currently under worldwide development by Novartis Crop Protection. In the course of development, the need for a practical and economically viable synthesis arose.

One synthesis of CGA 293'343 utilises the route shown in Fig 1. Reaction of 2-chloro-5-chloromethylthiazole **2** with the oxadiazinane intermediate **1** gave the desired product in good yield. Therefore, the core task to be addressed in the synthesis plan was the construction of the chlorothiazole moiety.

To date, apart from our work, five synthetic routes for compound **2** have been published. When we started out on our endeavour, only two references<sup>3,4</sup> were known. Neither route satisfied our need for a high-yielding, economically and ecologically sound process. After completion of our synthetic studies, further synthetic routes<sup>6–8</sup> were subsequently



**Figure 1.** Synthesis of CGA 293'343 from the intermediate **2**.

Reaction conditions: DMF, K<sub>2</sub>CO<sub>3</sub>, 2 h, 60°C, 74%.

\*Correspondence to: Thomas Pittner, Novartis Crop Protection AG, PO Box, CH-4002 Basel, Switzerland.  
E-mail: thomas.pittner@cp.novartis.com

(Received 30 June 1998; accepted 12 November 1998)

published, of which those in References 5 and 8 overlapped with our own, independent work.

## 2 SYNTHETIC ROUTES

**2.1 2-Chloro-5-chloromethylthiazole (2; Fig 1).** We followed two synthetic approaches, which are shown in Fig 2. Compounds 4, 5 and 6 are acyclic precursors with the functionality set up in such a way that, after cyclisation, the 2-chloro- and the 5-chloromethyl group are formed. In contrast, precursors 8 and 10 already possess the proper heterocyclic ring skeleton; conversion to 2 then proceeds in one step. Note that for compound 8 this means (i) replacement of the sulfur substituent by chlorine, and (ii) transformation of the exomethylene group into a chloromethyl group, both processes accompanied by a double bond shift leading to aromatisation. In the next section, another synthetic strategy will be discussed, in which the modifications (i) and (ii) are conducted in two separate steps.


The sodium salt 4 was prepared from the previously described amine 3 and carbon disulfide.<sup>9</sup> Oxidation with iodine or hydrogen peroxide gave the disulfide, which, on treatment with sulfonyl chloride, cyclised to give 2.<sup>10</sup> The isonitrile 6 was prepared from the corresponding formamide 5 by treatment with a dehydrating agent such as thionyl chloride ( $\text{SOCl}_2$ ). With  $\text{SCl}_2$ , the isonitrile 6 cyclised to form 2.<sup>11</sup> This process could also be performed in one

step by addition of  $\text{SOCl}_2$  and  $\text{SCl}_2$  to the formamide 5.


The thiazole 10, prepared from the commercially available aminothiazole 9 via a Sandmeyer reaction, could be halogenated on the methyl group with NCS to give 2.<sup>10</sup> Use of *N*-bromosuccinimide (NBS) gave 5-bromomethyl-2-chlorothiazole, which is also a useful intermediate for the synthesis of CGA 293'343. The previously described thiothiazole 8 was prepared in high yield from 7 and  $\text{CS}_2$ .<sup>12</sup> Treatment of 8 with chlorine or sulfonyl chloride gave 2.<sup>10</sup>

**2.2 The intermediate 2-benzylmercapto-5-chloromethylthiazole 12.** Synthesis of 2 from precursor 12 represents a third synthetic strategy, as shown in Fig. 3. Compound 12 has a chloromethyl group already in place, and the benzylmercapto substituent can be exchanged for chlorine. This can be done directly, leading to 2, or after attachment to the oxadiazinane intermediate 1, opening up an alternative reaction scheme as shown in Fig 4. As demonstrated in Fig 3, the intermediate 12 can be prepared by a variety of methods.


The commercially available mercaptothiazole 13 was alkylated to give 2-benzylmercaptothiazole 14. The intermediate 12 was prepared by treatment of 14 with DMF/ $\text{POCl}_3$ , subsequent catalytic hydrogenation of the product 15, and treatment of the resulting alcohol with  $\text{SOCl}_2$ .<sup>13</sup> 12 was then chlorin-



**Figure 2.** Synthesis of intermediate 2: a)  $\text{H}_2\text{O}$ ,  $\text{NaOH}$ ,  $\text{CS}_2$ , 85%, b)  $\text{H}_2\text{O}$ ,  $\text{KI}$ ,  $\text{I}_2$ , then  $\text{CH}_2\text{Cl}_2$ ,  $\text{SO}_2\text{Cl}_2$ , 30%, c)  $\text{HCO}_2\text{C}_2\text{H}_5$ , reflux, 12 h, 90%, d) DMF,  $\text{SOCl}_2$ ,  $\text{Na}_2\text{CO}_3$ , 12 h, 80%, e)  $\text{CCl}_4$ ,  $\text{SCl}_2$ ,  $40^\circ\text{C}$ , 4 h, 50%, f)  $\text{SOCl}_2$ ,  $\text{SCl}_2$ , reflux, 24 h, 42%, g) *t*-BuONO,  $\text{CuCl}_2$ ,  $\text{CH}_3\text{CN}$ , 60%, h) *N*-chlorosuccinimide, dibenzoylperoxide,  $\text{CCl}_4$ , reflux, 50%, i)  $\text{CS}_2$ , EtOH, ambient temperature, 2 h, 90%, j)  $\text{CH}_2\text{Cl}_2$ ,  $\text{H}_2\text{O}$ ,  $\text{SO}_2\text{Cl}_2$ , 1 h, 80%.



**Figure 3.** Synthesis via the intermediate 12: a)  $\text{CH}_3\text{CN}$ ,  $\text{K}_2\text{CO}_3$ , benzylbromide,  $60^\circ\text{C}$ , 1 h, 70%, b)  $\text{CH}_2\text{Cl}_2$ ,  $\text{NaHCO}_3$ ,  $\text{SO}_2\text{Cl}_2$ , 60%, c)  $\text{CH}_2\text{Cl}_2$ , 2 h, 60%, d)  $\text{CH}_3\text{CN}$ ,  $\text{K}_2\text{CO}_3$ , benzylbromide,  $60^\circ\text{C}$ , 1 h, 80%, e) DMF,  $\text{POCl}_3$ ,  $40^\circ\text{C}$ , 6 h, 35%, f)  $\text{H}_2$ ,  $\text{PtO}_2$ ,  $\text{FeCl}_2 \cdot 4 \text{H}_2\text{O}$ , ethyl acetate, then  $\text{SOCl}_2$ ,  $\text{CH}_2\text{Cl}_2$ , 3 h, 60%, g) ethanol,  $\text{H}_2\text{O}$ , glycidyl aldehyde, 12 h, 85%, h)  $\text{CH}_2\text{Cl}_2$ ,  $\text{SOCl}_2$ , 3 h, 90%.



**Figure 4.** Alternative synthesis of CGA 293'343 from the intermediate **12**. Reaction conditions: a) oxadiazinane **1**, DMF,  $K_2CO_3$ , 2 h, 60°C, 75%, b) HCl,  $H_2O$ ,  $C_6H_5Cl$ ,  $Cl_2$ , 6 h, 80%.

ated to give **2**. Another synthetic approach to **12** is chlorination of 2-benzylmercapto-5-methylene-thiazoline **11**,<sup>14</sup> which is available by alkylation of **8** with benzylbromide. **12** was also prepared by treatment of diol **17** with  $SOCl_2$ .<sup>13</sup> Diol **17** was constructed from the previously described dithiocarbamate **16** by reaction with glycidyl aldehyde.<sup>15</sup>

An alternative synthesis is shown in Fig 4. Coupling of **12** with **1** followed by chlorination of the intermediate **18** gave CGA 293'343.<sup>13</sup>

## REFERENCES

- 1 Maienfisch P and Gsell L (Ciba-Geigy AG), Preparation of 3-(heterocyclymethyl)-4-iminoperhydro-1,3,5-oxadiazine derivatives as pesticides. *Eur Pat Appl* EP 580553 A2 940126 (1994).
- 2 Yamamoto I, *Agrochem Jpn* **68**:14 (1996).
- 3 Beck G and Heitzer H (Bayer AG), Manufacture of 2-chloro-5-(chloromethyl)-thiazole. DE 3631538 A1 880324 (1988).
- 4 Uneme H, Higuchi N and Minamida I (Takeda Chem. Ind Ltd), Preparation of chlorthiazole derivatives. *EP* 446913 A1 910918 (1991).
- 5 Kraatz U (Bayer AG), Preparation of 2-chloro-5-chloromethyl thiazole. *EP* 780384 A2 970625 (1997).
- 6 Matsuda H, Asanuma G and Shiono M (Kuraray Co Ltd), Preparation of 2-chloro-5-chloromethyl thiazole by treatment of 3-chloro-1-isothiocyanato-1-propene with a chlorinating agent. *EP* 794180 A1 970910 (1997).
- 7 Jackson A, Heyes G, Greyson JI and Clarke R (Fine Organics Ltd Agro-Kanesho Ltd), Preparation of 2-halo-5-haloalkyl thiazoles. *EP* 763531 A1 970319 (1997).
- 8 Wakasugi T, Miyakawa T and Tanonaka T (Kureha Chem Ind Co Ltd), *EP* 775700 A1 970528.
- 9 Bargar M *et al.*, *J. Org Chem* **52**:678-681 (1987).
- 10 O'Sullivan AC, Gsell L, Naef R, Senn M, Pitterna T and Wadsworth DJ (Ciba-Geigy AG), Preparation of 2-chloro-5-chloromethyl thiazole. WO-A-97/23469 (1997).
- 11 Pitterna T (Ciba-Geigy AG), Preparation of 2-chloro-5-methyl thiazole compounds. WO-A-97/10226 (1997).
- 12 Hanefeld W and Bercin E, *Liebigs Ann Chem* **1**:58-64 (1985).
- 13 Pitterna T, Szczepanski H, Maienfisch P, Hüter OF, Rapold T, Senn M, Göbel T and O'Sullivan AC (Ciba-Geigy AG), Preparation of thiazoles. WO 9827075 A1 980625 (1998).
- 14 Szczepanski H, Göbel T, Hüter OF, O'Sullivan AC, Senn M, Rapold T, Maienfisch P and Pitterna T (Ciba-Geigy AG), Preparation of 2-chloro-thiazoles. WO 9720829 A1 970612 (1997).
- 15 von Braun J, *Chem Berichte* **35**: 3368 (1902).

## Non-steroidal ecdysone agonists: New tools for IPM and insect resistance management

Tarlochan S Dhadialla\* and Richard K Jansson  
Insecticide Discovery Group, Rohm and Haas Company,  
727 Norristown Road, Spring House, PA 19477, USA

**Abstract:** The non-steroidal, bis-acylhydrazine agonists of the insect molting hormone, 20-hydroxyecdysone, were first discovered over ten years ago. An extensive structure-activity optimization program yielded one commercial insecticide, tebufenozide (RH-5992) and two additional candidate insecticides (methoxyfenozide and halofenozide) which are in development. Tebufenozide is highly selective for lepidopteran pest control and is thus useful for IPM and resistance management programs. Methoxyfenozide (RH-2485) is also lepidopteran-selective but significantly more potent than tebufenozide and offers control of a wider range of lepidopteran pests. Halofenozide (RH-0345) is generally less potent and selective than tebufenozide or methoxyfenozide. However, its physical and biological properties make it well suited for control of beetle grubs and caterpillars in the soil. Target pest selectivity, new and novel mode of action, ecotoxicological safety and safety to beneficial arthropods make these insecticides valuable tools for integrated pest and resistance management programs.

© 1999 Society of Chemical Industry

**Keywords:** tebufenozide; methoxyfenozide; halofenozide; ecdysone agonists; RH-5992; RH-2485; RH-0345; IPM; resistance management programs

The steroid insect moulting hormone, 20-hydroxyecdysone (20E), and the sesquiterpenoid juvenile hormone play a central role in the regulation of the growth and development, as well as of reproductive processes, in insects. As such, chemicals which mimic or antagonize the action of these two hormones have been sought for use as safe, third-generation pesticides. While success in the discovery of juvenile hormone mimetics came much earlier,<sup>1</sup> it is only recently that insecticides which act as agonists of 20E have been discovered.<sup>2</sup>

Scientists at Rohm and Haas Company have discovered three non-steroidal ecdysone agonists, all of which belong to bis-acylhydrazine chemistry.<sup>2</sup> One of these, *N*-tert-butyl-*N'*-(4-ethylbenzoyl)-3,5-dimethylbenzohydrazide (tebufenozide; RH5992) was the first to be commercialized as a lepidopteran-specific insecticide under the trade names Mimic®, Confirm® and Romdan® in several countries.

\* Correspondence to: Tarlochan S Dhadialla, Rohm and Haas Co, 727 Norristown Road, Spring House, PA 19477, USA.  
E-mail: RSATSD@rohmmaas.com  
(Received 1 July 1998; accepted 12 November 1998)